


Binomial

Binomial(p,n)

[Binomial Equations](#)

[Crystal Ball parameter restrictions](#)

A Binomial(p,n) distribution returns discrete values between 0 and n. Examples of the Binomial distribution are shown below:



Uses

The Binomial distribution models the number of successes from n independent trials where there is a probability p of success in each trial (as explained in the section on the [Binomial process](#)).

The binomial distribution has an enormous number of uses. Beyond simple [binomial processes](#), many other stochastic processes can be usefully reduced to a binomial process to resolve problems. For example:

Binomial process:

- Number of false starts of a car in n attempts;
- Number of faulty items in n from a production line;
- Number of n randomly selected people with some characteristic;

Reduced to binomial:

- Number of machines that last longer than T hours of operation without failure;
- Blood samples that have zero, or >0 antibodies;
- [Approximation](#) to a hypergeometric distribution

The following links lead to just some of the examples and models in ModelAssist that use the binomial distribution:

[Conditional logic](#)

[Sampling from a liquid](#)

[Distribution fitting of threshold data](#)

[Bayesian prior](#)

[Test result](#)

Comments

The Binomial distribution makes the assumption that the probability p does not change the more trials are performed. That would imply that my aim doesn't get better or worse. It wouldn't be a good estimator, for instance, if the chance of success improved with the number of trials.

Another example: the number of faulty computer chips in a 2000 volume batch where there is a 2% probability that any one chip is faulty = Binomial (2%, 2000).

The Binomial distribution was first discussed by Bernoulli (1713). It is related to the [Beta](#) and [Negative Binomial](#) distributions, all of which have their basis in the [Binomial process](#) where the Binomial distribution is also derived. The [Bernoulli](#) distribution is a special case of the Binomial with $n = 1$ i.e.: Bernoulli (p) = Binomial($p, 1$) that is used to model risk events.

The Binomial distribution has the property $\text{Binomial}(p, n) + \text{Binomial}(p, m) = \text{Binomial}(p, n+m)$ which makes sense if one thinks of n and m being two sets of independent binomial trials, all with the same probability of success.

The Excel function `BINOMDIST(s,n,p,0)` returns the binomial probability mass function, and `BINOMDIST(s,n,p,1)` returns the binomial cumulative distribution function.
